

<https://helda.helsinki.fi>

Role of Biochar and Fungi on PAH Sorption to Soil Rich in Organic Matter

Anasonye, Festus

2018-02-01

Anasonye , F , Tammeorg , P , Parshintsev , E , Riekkola , M-L & Tuomela , M T 2018 , ' Role of Biochar and Fungi on PAH Sorption to Soil Rich in Organic Matter ' , Water, Air and Soil Pollution , vol. 229 , no. 2 , 49 . <https://doi.org/10.1007/s11270-018-3708-2>

<http://hdl.handle.net/10138/307851>

<https://doi.org/10.1007/s11270-018-3708-2>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Role of biochar and fungi on PAH sorption to soil rich in organic matter**

2

3 Festus Anasonye^{a,*}, Priit Tammeorg^b, Jevgeni Parshintsev^c, Marja-Liisa Riekkola^c, Marja Tuomela^a

4 ^aDepartment of Food and Environmental Sciences, P.O. Box 56, Biocenter 1, FI-00014 University of Helsinki, Finland

5 ^bDepartment of Agricultural Sciences, P.O. Box 27, Plant Production Sciences, FI-00014 University of Helsinki, Finland

6 ^cLaboratory of Analytical Chemistry, Department of Chemistry, A.I. Virtasen aukio 1 (P.O. Box 55) FI-00014 University
7 of Helsinki, Finland

8 *Corresponding author's e-mail address: anasonyefestus@yahoo.com

9

10 Key words: ACTIVATED BIOCHAR, AMENDMENT, HAPLIC ARENOSOL, ORGANIC MATTER, PYRENE,
11 SORPTION

12 **ABSTRACT**

13 The use of biochar (BC) has been suggested for remediation of contaminated soils. This study aims to investigate the role
14 of microorganisms in sorption of PAH to BC-amended soils. Fungi, especially the wood and litter-degrading fungi, have
15 shown the ability for humification and to degrade recalcitrant molecules, and are thus suitable model organisms. Haplic
16 Arenosol with high organic matter content was chosen to highlight the importance of soil organic matter (SOM) in PAH
17 sorption, possibly to form non-extractable residue. Basidiomycetous fungi *Agrocybe praecox* and *Phanerochaete*
18 *velutina* grown on pine bark were inoculated in organic matter (OM) rich Haplic Arenosol and OM poor sandy loam with
19 either BC or chemically activated BC (ABC) and ¹⁴C- labelled pyrene for 60 days. Fungi did not mineralize pyrene, but
20 increased sorption up to 47–56% in BC-amended Haplic Arenosol in comparison with controls (13–25%) without a
21 fungus irrespective of the presence of an adsorbent. In OM poor sandy loam only 9–12% of pyrene was sorbed to amended
22 soil in the presence of fungus and adsorbent. Combining BC and fungi is an effective method for sorbing pyrene especially
23 in high SOM soils.

24 **1 Introduction**

25 Polyaromatic hydrocarbons (PAHs) are organic contaminants widespread in the environment in many parts of the world.
26 In Europe alone, PAHs contamination accounts for 11% of 340,000 contaminated sites (Liedekerke et al. 2014). A major
27 source of PAHs contamination in soil is creosote, a coal-tar distillate used as a preservative for power lines and crossties
28 (Murphy and Brown 2005). PAHs are also formed by incomplete combustion of biomass. PAHs persist in the environment
29 due their low water solubility, complex chemical structure and reduced degradation (Winquist et al. 2014; Lamichhane
30 et al. 2016). Nearly all PAHs are highly toxic; thus, they are of concern to all life forms in soil (Chen and Liao 2006).
31 Bioavailability of PAHs in soil is governed by soil organic matter (SOM) content and presence of naturally occurring or
32 added adsorbents (Macleod and Semple 2002; Cornelissen et al. 2006; Lamichhane et al. 2016). The fate of PAHs in soil

depends on many physical, chemical and biological processes such as absorption, volatilization, photolysis, chemical degradation, and microbial degradation (Deng and Zeng 2017).

Biochar (BC) is produced from biomass by pyrolysis and is distinguished from other carbonaceous materials by its end use (EBC 2012). Differently from charcoal (often used for energy production), BC is used in a way that does not involve rapid mineralisation of the photosynthetically fixed carbon back to atmosphere (EBC 2012). BC has shown capacity to sorb organic contaminants (Beesley et al. 2011; Zhu et al. 2017), however, its sorption capacity is lower compared with activated carbon (AC; Hale et al. 2011). Apart from contaminant sorption, application of BC to soil offers other advantages, such as carbon sequestration (Woolf et al. 2010; Smith, 2016), soil fertility improvement (Major et al. 2010; Jones et al. 2012; Tammeorg et al. 2014a; b; Ding et al. 2016) and reduction of N₂O emissions from soil (Case et al. 2012; Angst et al. 2013; Zhu et al. 2017). Thus, BC is a relevant alternative to AC despite its moderate sorption capacity.

The influence of BC addition on soil microorganisms have been increasingly studied (e.g. Pietikäinen et al. 2000; Lehmann et al. 2011; Abujabhah et al. 2016; Dai et al. 2016; 2017). Pores in BC can serve as home for microorganisms (Warnock et al. 2007; Quilliam et al. 2013a) and also harbour microorganisms that are not native to soils, while surfaces can provide a platform for biofilm formation (Lehmann et al. 2011; Noyce et al. 2016). Changes in microbial composition and abundance in soil following BC addition have been reported (Pietikäinen et al. 2000; Nielsen et al. 2014; Mitchell et al. 2016; Abujabhah et al. 2016; Dai et al. 2017). O'Neill et al. (2009), Grossman et al. (2010) and Taketani et al. (2013) reported higher microbial population and an increased microbial diversity in high native black carbon Anthrosols than in adjacent soils. Increase in relative abundance of soil bacteria and a decrease in soil fungi with BC addition has been reported and the changes was due to an increase in soil carbon content as BC may have supplied labile C substrates that favored fast growing bacteria over fungi (Khadem and Raiesi 2017). BC addition stimulates biological processes such as increases in enzyme activities and respiration rates in amended soil (AlMarzooqi and Yousef 2017). Jin (2010) observed a shift in a fungal community following BC amendment. The observed changes in microbial communities following BC amendment may also be explained by increase in nutrient availability and utilization which leads to an overall increase in soil fertility (Kolton et al. 2011; Anderson et al. 2011; Nielsen et al. 2014; Pan et al. 2016). Other key factors controlling the shifts in relative abundance and diversity of microbial populations in soil following biochar addition may include; soil physicochemical properties such as organic matter content, pH and texture, type of biochar applied, incubation time and climatic conditions (Lehmann et al. 2011; Farrel et al. 2013; Prayogo et al. 2013; Ogonnaya et al. 2014; Watzinger et al. 2014; Zhu et al. 2017).

Although BC contributes greatly in sorption of PAH compounds in the BC-amended soil (Zhang et al. 2010; Chen and Yuan 2011), very little is known about the role of microorganisms especially the fungi in this sorption process (Quilliam et al. 2013b; Zhu et al. 2017). PAHs in BC-amended soil can be degraded by both bacteria and fungi in soil (Rhodes et al. 2008; Quilliam et al. 2013b) as BC can enhance action of fungi in soil (García-Delgado et al. 2015). PAHs and other aromatic compounds in the amended soil can be oxidized and incorporated into SOM as non-extractable or bound residues by enzymes produced by fungi in the soil (Dec et al. 2001; Kästner et al. 2014). The processes of enhanced sorption by BC and enhanced formation of non-extractable or bound residues from PAH compounds by fungi in soil have been adopted as effective soil remediation strategies as they reduce the bioavailability of PAH in soil (Bollag 1992; Kästner et al. 2014; Zhu et al. 2017). The possibility that both strategies, processes of enhanced sorption by BC and enhanced formation of non-extractable or bound residues from PAH compounds by fungi occurring in soil amended with BC has

not been previously studied. PAH degradation in soil by fungi as well as sorption to BC-amended soil differs between soil types (Anyika et al. 2015). The fate of PAH in BC-amended soil may depend on the properties of the soil and BC as well as microbial composition of the amended soil (Anyika et al. 2015).

In this study, we investigated the role of microorganisms in sorption of PAH to BC-amended soils when combined with fungi. Pyrene, a common pollutant in PAH-contaminated soil that is not readily degraded by other microorganisms in soil was selected as a model PAH compound. The aim was to understand the role of fungi, BC and their combined effect on the environmental fate of PAH in contaminated soil using ^{14}C -pyrene-labelled compound. A particular emphasis was on the role of soil organic matter, as almost all the previous studies have been performed with OM poor soil. We hypothesized that a) the porous carbonaceous soil amendments will sorb the more PAH the higher their specific surface area and b) that PAH are more easily sorbed to biochar in low SOM soils and c) that introducing fungi to the system will further increase the sorption of PAH to high SOM soil by oxidative enzymes binding organic compounds to SOM.

2 Materials and methods

2.1 Chemicals

PAH mix, containing 16 United States Environmental Protection Agency (US-EPA) PAH compounds (acenaphthene, acenaphthylene, anthracene (ANT), benzo(a)anthracene, benzo(a)pyrene (BaP), benzo(b)fluoranthene, benzo(g,h,i)perylene, benzo(k)fluoranthene, chrysene, dibenz(a,h)anthracene (DBA), fluoranthene, fluorene, indeno(1,2,3-cd)pyrene, naphthalene, phenanthrene, pyrene (PYR), and carbazole at each concentration of 2 mg mL^{-1} in dichloromethane : benzene (1:1) was purchased from AccuStandard, New Haven, CT, USA. Sodium acetate (American Chemical Society certified grade), sodium chloride (analytical grade), isopropanol (analytical grade) and toluene (HPLC grade) were purchased from Sigma-Aldrich (St. Louis, MO, USA). 1,1'-binaphthyl (97% purity) used as the internal standard during PAH analysis was purchased from Acros Organics, (Geel, Belgium). 4,5,9,10- ^{14}C -pyrene (specific activity $2.035\text{ GBq mmol}^{-1}$, radiochemical purity $\geq 99\%$) was purchased from Sigma-Aldrich (St. Louis, MO, USA).

2.2 Soils and adsorbents used in the experiment

Three non-contaminated soils were used in the experiments with BC and one PAH contaminated soil was used for PAH extraction method optimization and screening of fungi. The first soil has a texture class of sandy loam and was sampled from a depth of 0–20 cm (a non-agricultural soil) from Biocenter 1 surroundings of Viikki campus of University of Helsinki, Helsinki Finland ($60^{\circ}22'\text{N}$, $25^{\circ}01'\text{E}$). The second soil has a texture class of sand and was sampled from top 0–20 cm depth (an agricultural soil) from Viikki agricultural field, University of Helsinki, Helsinki Finland ($60^{\circ}22'\text{N}$, $25^{\circ}02'\text{E}$). The third soil is classified as an Entic Haplocryod (Soil Survey Staff, 1998) or a Haplic Arenosol (FAO-UNESCO, 1997) and is simply referred here as Haplic Arenosol, was collected in the vicinity of Forest Field Station of the University of Helsinki in Juupajoki, ($61^{\circ}84'\text{N}$, $24^{\circ}26'\text{E}$) (corresponds to Hyytiälä soil in Ilvesniemi et al. 2000). Soil samples were stored at 4°C before use. PAH contaminated soil which is described in detail in Winquist et al. (2014), was used to screen for fungal growth on BC-amended contaminated soil and for PAH extraction optimization.

BC was prepared from spruce wood chips as described in Tammeorg et al. (2014a). AC used was a commercial product from coconut shell charcoal (AC004; Activated Carbon Technology UK Limited, Billingham, UK). Particle size of AC

106 was 2.36–4.75 mm. ABC was produced from BC (see the next section). Prior to the experiment with soil, BC and ABC
107 were sieved with < 2 mm mesh. AC was used as received from the producer.

108 pH of adsorbents, soil and soil amended with adsorbents was measured from 1:2.5 (v v⁻¹) suspension in deionized water,
109 while moisture contents were measured gravimetrically by drying a known mass of soil overnight at 105°C. Soil organic
110 matter content (SOM) was measured by placing dried soil at 550°C for 4 hours and calculated as loss on ignition. The
111 specific surface area of BC, ABC and AC was measured by the Brunauer, Emmett, and Teller (BET) nitrogen adsorption
112 method measured with Micromeritics surface analyzer (Micromeritics Tristar II 3020, Norcross, GA, USA) at Tampere
113 University of Technology, Finland. The elemental carbon, hydrogen and nitrogen content of samples was measured by
114 using a VarioMax elemental analyzer (Elemental Analysensysteme GmbH, Hanau, Germany). Cation exchange capacity
115 (CEC) of adsorbents and soil was analyzed by adopting a method described by Mitchell et al. (2013) with some
116 modifications. A sample of 0.5 g of adsorbent or 1 g of soil with or without adsorbent was used. The solution was analyzed
117 using flame photometer (Model 410, Corning, New York, USA). The contribution of native black carbon to total organic
118 carbon content of the soil was estimated using the peroxide/weak nitric acid digestions method described by Kurth et al.
119 (2006), except that soil was ground to 0.63 µm particle size. The total carbon remaining, estimated as oxidation-resistant
120 elemental carbon was considered to be the soil native black carbon (Rumpel et al. 2006).

121 2.3 Activated biochar production

122 Activated biochar (ABC) was produced from BC as described by Lalhruaitluanga et al. (2011). In brief, sieved BC was
123 added to potassium hydroxide (KOH) solution of various concentrations (1, 10, 20, 40, 60 and 80%). The mixture was
124 stirred for 24 h at a speed of 150 rpm. ABC was separated from KOH solution by centrifugation at 10,000 rpm for 10 min
125 and thoroughly washed with deionized water, to remove excess KOH and other impurities that might be blocking the
126 newly created pores on the surface of the activated biochar. The separated ABC was oven-dried at 110 °C for 24 h and
127 then cooled at room temperature and stored in an air-tight container.

128 2.4 PAH sorption to soil and soil amended with adsorbents

129 To prepare the soil with desirable amendments, i.e. AC and ABC (1%), and BC (1 and 2%) (w/w) samples, soil was
130 thoroughly mixed with an adsorbent. For each treatment, 10 g of soil and either AC, BC or ABC was added to a 100 mL
131 glass bottle. Soil-adsorbent mixture in glass bottle was spiked with 125 µL of 2 g L⁻¹ mixture of four PAHs, comprising
132 ANT (3-rings), PYR (4-rings), BaP (5-rings) and DBA (5-rings) in dichloromethane benzene (1:1). Acetone (10 mL) was
133 added to the mixture and mixed thoroughly to ensure homogeneous mixing of PAHs in soil. After mixing, acetone was
134 allowed to evaporate in the hood for 3 days. The final concentration for each PAH in the soil was 25 mg kg⁻¹ (i.e. total
135 PAH concentration in the soil was 100 mg kg⁻¹). Soils without adsorbents were spiked with the same concentration of
136 PAHs in order to obtain control samples. The first bottles were taken for extraction immediately after acetone had
137 evaporated, i.e. on day 3. The remaining bottles were incubated at 21°C in the dark for 60 days.

138 2.5 PAH extraction

139 PAH concentration of adsorbents was measured after Soxhlet extraction as described in Tammeorg et al. (2014a). Three
140 PAH extraction methods were evaluated: ultrasound-assisted extraction (UAE), accelerated solvent extraction (ASE) and
141 Soxhlet extraction (SE) for contaminated soil and details are in Fig. S1. For further analyses, ASE method was selected

142 as it was automatic, efficient, fast, easy and environmentally safe, and PAHs were extracted as efficiently as with Soxhlet
143 which is considered a conventional extraction method. ASE condition used are as that described by Hilber et al. (2012).

144 2.6 PAH analysis by gas chromatography-mass spectrometry (GC-MS)

145 PAHs were analysed with gas chromatography (Agilent Technologies 6890N, USA) equipped with mass spectrometry
146 (MS Agilent 5973N) by injecting 1 μL of the extract using splitless injection mode. Separation was done on Zebron High
147 performance ZB-5ms GC column (30 m, 0.25 mm internal diameter, 0.25 μm film thickness) from Phenomenex, USA.
148 A deactivated retention gap (Agilent Technologies) of 2 m length with 0.53 mm internal diameter was placed before the
149 separation column. Helium was used as a carrier gas at constant pressure of 100 kPa. The injection temperature was set
150 to 320°C while the oven temperature program was as follows: 90°C (2 min); increase of 10°C min⁻¹ until 320°C; 320°C
151 (15 min). Detection was performed with MS in the electron impact mode with 70 eV ionization energy. The ion source
152 temperature was 150°C and the interface between the GC and the quadrupole MS was set to 320°C. Compounds were
153 identified with the National Institute of Standard and Technology (NIST) library and quantification was done using
154 internal standard method. Toluene mixtures containing varying amounts of analytes (concentrations = 0.1, 0.5, 1.0, 2.0,
155 4.0 $\mu\text{g mL}^{-1}$) and a constant amount of internal standard (1,1'-binaphthyl 5 $\mu\text{g mL}^{-1}$) were used for calibration. The limits
156 of detection for ANT, PYR, BaP and DBA were 0.02, 0.03, 0.04 and 0.06 $\mu\text{g mL}^{-1}$, respectively.

157 2.7 Sorption experiment

158 The sorption coefficient (K_d) of ¹⁴C pyrene was obtained in a batch equilibration experiment with three replicates as
159 described by Kumari et al. (2014) with little modifications. Three millilitres from the soil solution was taken and mixed
160 with 10 mL of scintillation cocktail (OptiPhase "HiSafe" 3® Perkin Elmer Inc.; Fisher Chemicals, Loughborough
161 Leicestershire, England). The radioactivity was measured using a liquid scintillation counter (Wallac 1411, Scintillation
162 Products, Wallac Oy, Turku, Finland). BC and ABC-amended soils, (sandy loam, sand and Haplic Arenosol) were used
163 in sorption experiment with two variants; 1) non-extracted soil and 2) DOC-extracted soil. DOC was extracted from soil
164 as described by Impellitteri et al. (2002) except that here 3 further rounds of extractions were carried out.

165 2.8 Fungal strains, inoculation and screening

166 Fungal strains used in this experiment were obtained from the Fungal Biotechnology Culture Collection (FBCC) of the
167 Department of Food and Environmental Sciences, University of Helsinki, Helsinki, Finland, and were grown on 2% malt
168 extract agar plates and incubated at 25 °C before the use in experiments. Basidiomycetous fungi were chosen as model
169 organisms due to their unique ability to decontaminate PAH contaminated soil by mineralization and humification of
170 PAH compounds. The fungal species with FBCC number (isolation number is given in parenthesis) were: *Agrocybe dura*
171 478 (Mn71-2), *Agrocybe praecox* 476 (Tm70.84), *Phanerochaete velutina* 941 (T244i), *Obba rivulosa* 939 (T241i)
172 (formerly known as *Physisporinus rivulosus*), *Rhodocollybia butyracea* 626 (K209) and *Stropharia coronilla* 480 (stock
173 gram B). Fungal liquid cultures were prepared according to Anasonye et al. (2015). Fungi were incubated on autoclaved
174 Scots pine (*Pinus sylvestris*) bark, until the surface of the bark was fully covered with fungal mycelium. The screening of
175 fungal strains was performed in PAH contaminated soil amended with either 1% or 2% (w/w) BC. Petri-dishes of 9 cm
176 in diameter were filled with 50 g of soil and 2.5 g pine bark with fungal mycelium was placed on top of the soil. Fungal
177 growth was examined visually once a week during 90 days of incubation at room temperature in the dark. Moisture
178 content of the soil was kept constant by adding water if needed.

179 2.9 Sorption of ¹⁴C-pyrene

180 Two fungi were selected for further experiments, namely *A. praecox* and *P. velutina*. Non-sterile sandy loam or Haplic
181 Arenosol (10 g) was placed in a 100 mL glass bottle (Schott Duran laboratory glassware, Mainz, Germany) and ¹⁴C-
182 labelled pyrene was added uniformly to soil (radioactivity 2,100 Bq per bottle) and thoroughly mixed to ensure
183 homogeneous result. Sand was not used in this experiment as two soil types with marked difference in SOM was preferred.
184 The soil was amended with either BC or ABC, 1% for sandy loam and 2% for Haplic Arenosol (w/w). Soil moisture
185 content was adjusted to 40% of maximum water holding capacity using deionized water and 5 g of fungal inoculum
186 growing on bark was added on top of the soil. For control bottles, uninoculated bark was added on top of spiked soil.

187 Abiotic control bottles were included to account for the impact of indigenous microorganisms on pyrene sorption and
188 degradation. For abiotic control, soil was autoclaved before spiking with ¹⁴C-pyrene and adding adsorbent. All treatments
189 with three replicates were incubated at 21°C in the dark for 60 days. The bottles were flushed with moist air for 15 minutes
190 once a week. Mineralized fraction, i.e. evolved ¹⁴CO₂ was trapped during the aeration period into 10 mL of 1 M NaOH.
191 Radioactivity was measured from 1 mL of trapping solution mixed with 10 mL of scintillation cocktail (OptiPhase
192 “HiSafe”). After incubation, bark with or without fungus was separated from soil by sieving. Pyrene was extracted from
193 the soil with ASE as described earlier. The radioactivity was measured with liquid scintillation counter from 1 mL of
194 toluene extract mixed with 10 mL of the scintillation cocktail. After toluene extraction, soil and sieved bark were
195 combusted separately with Junitek Oxidizer (Junitek Oy, Turku, Finland). During combustion, all the carbon in the sample
196 forms CO₂, which is trapped with a mixture of 16 mL Lumasorb2 II and Carboluma2 scintillation liquids (1:1 v/v; Lumac
197 LSC, Belgium) and radioactivity is measured with liquid scintillation counter. The mass balance of ¹⁴C-pyrene was
198 determined as in Valentin et al. (2013) with modifications: (1) toluene extraction (available fraction), (2) fraction bound
199 to soil (unavailable fraction), and (3) fraction bound to bark.

200 2.10 Statistical analyses

201 The effects of experimental treatment combinations on the percentage of PAH sorbed were tested with a two-way analysis
202 of variance (ANOVA) with soil type, treatment, and their interactions as fixed effects for both incubation times (3 and 60
203 days). The effects of experimental treatment combinations on the K_d and mass balance were tested with a two and three-
204 way analysis of variance (ANOVA), respectively with soil type, treatment, and their interactions as fixed. Means were
205 compared using the Tukey HSD multiple pair-wise comparison test at p < 0.05. The normal distribution of the residuals
206 from the models was tested with Shapiro-Wilk test and the homogeneity of variances was tested with Levene's test. If the
207 data failed to meet the assumptions for parametric statistics, Box-Cox transformation was used (Box and Cox, 1964).
208 Statistical analyses were carried out with the software package PASW v 20.0 (SPSS Corp., Chicago, USA).

209 3 Results

210 3.1 Physicochemical properties of soils and adsorbents

211 Haplic Arenosol had lower pH and significantly higher DOC and SOM contents compared with sandy loam and sand
212 (Table 1). The highest content of native black carbon was recorded in Haplic Arenosol (2% of soil total mass) while the
213 lowest was in sandy loam soil. Elemental compositions of BC, ABC and AC were similar, but specific surface area of

AC was considerably higher than that of ABC and BC (Table 2). Total PAH (US-EPA 16 PAH compound) concentration of adsorbents were below or within the recommended maximum limit set by international BC initiative (6–30 $\mu\text{g g}^{-1}$; IBI 2015) and below the limit by European Union (12 $\mu\text{g g}^{-1}$; EBC 2012) for use as a soil amendment in agriculture.

217

218 3.2 Sorption capacity of BC

Soil types, treatments and interactions affected the percentage of PAHs sorbed for all four PAHs used in this experiment (Table 3). ABC-amended sandy loam sorbed (95–100%) PAHs in 3 days (Table S1). However, the effect seemed to be only short-lived as the sorption efficiency of ABC had decreased notably after 60 days of incubation. In sandy loam significantly, more pyrene was sorbed than in other soil types. AC addition to sandy loam and sand significantly enhanced pyrene sorption ($p < 0.001$, Table 3, Table S1) compared to BC and ABC during a 60-day incubation period.

224

225 3.3 Role of dissolved organic carbon (DOC)

BC and ABC addition significantly (Figure 1, $p < 0.001$, Table S2) increased ^{14}C -pyrene sorption to both sandy loam and sandy soils and the increase with ABC in both soils were larger compared with BC. In Haplic Arenosol, BC and ABC addition had little or no effect on pyrene sorption. When DOC was removed from the soil, pyrene sorption to control sandy loam and sand without amendments significantly (Fig 1, $p < 0.001$) decreased although it significantly improved in Haplic Arenosol. BC and ABC additions to DOC-extracted soil significantly (Fig 1, $p < 0.001$, Table S2) increased pyrene sorption to sandy loam soil.

232

233 3.4 Fungal treatment

All the screened fungi grew to the soil amended with BC except *Obba rivulosa* that grew only on bark used as a carrier material in inoculation (data not shown). Regardless of whether BC was added at 1 or 2 %, (w/w) there was no difference in growth of fungi. However, two fungi, *A. praecox* and *P. velutina* were chosen for further experiment, as they have shown in previous studies to have the capability to colonize the soil, compete favorably with indigenous microorganisms and to degrade PAH in soil (Steffen et al. 2002; Winqvist et al. 2014).

There was significant difference in the levels of pyrene that was sorbed to the soils used in the study ($p < 0.001$, Table 4). The role of fungal action to pyrene sorption in BC or ABC-amended soils was evaluated in two soil types. Higher levels of ^{14}C -pyrene was sorbed to Haplic Arenosol incubated with fungus (47–56%) compared with sandy loam (up to 19%). The lowest levels of pyrene sorption in the soil were observed in autoclaved soils with BC or ABC (9–15%), highlighting the role of indigenous microorganisms in PAH sorption to soil. Three-way analysis of variance (ANOVA) also showed that soil type significantly influenced pyrene sorption to the bark used in the experiment ($p < 0.001$, Table 4). Significantly higher levels of pyrene sorbed to sandy loam (27–36%) than to Haplic Arenosol (2–7%). There was no significant difference between the two fungal strains used in the way they caused change in sorption of pyrene to BC-amended soil. Both fungi significantly increased the sorption of pyrene to Haplic Arenosol-BC complex (Table 4). Similarly, the two

adsorbents used had no significant difference in the way they cause changes in sorption of pyrene to soil incubated with fungi. Haplic Arenosol amended with either of the adsorbents showed increased capacity to sorb pyrene compared to OM poor soil.. Haplic There was no evolution of $^{14}\text{CO}_2$ in either control or treated soil during the entire incubation period.

4 Discussion

PAH sorption was maximum mostly with ABC in the beginning of incubation regardless of soil type. This is particularly interesting considering that AC had almost double BET surface area compared with ABC or BC, thus our first hypothesis was only partially supported by the evidence. The phenomenon could be explained by the formation of alkaline surface sites on BC during activation with potassium hydroxide or wider pore size of ABC compared to AC and BC (de Andrés et al. 2013). Chemically activated char is known to form surface sites leading to enhanced sorption capacity (de Andrés et al. 2013; Lamichhane et al. 2016) and less time is needed for PAH to access sorption sites in ABC than AC or BC, if pore size in ABC is wider. Also, alkali modified BC enhances BC sorption capacity through the formation of π - π electron donor acceptor interaction between modified BC and aromatic rings of PAH (Liu et al. 2012; Zhu et al. 2017). Hence, BET surface area is not the only factor controlling PAH sorption, but other chemical and physical properties as well as the aliphatic regions of adsorbents play a role in contaminant sorption (Park et al. 2013). However, when ABC was aged for 60 days in soil, it was unable to retain most of the sorbed PAH. During a prolonged incubation in soil, the surface chemistry and sorption characteristics of ABC may have been altered (Cheng and Lehmann 2009; Gibson et al. 2016), resulting in the release of already sorbed PAH. This phenomenon is typical to char produced by chemical activation with potassium hydroxide and such effect was not detected with AC and BC. In a field scale experiment, Martin et al. (2012) observed release of already sorbed diuron and atrazine from BC in soil during aging, but when BC dose was increased, sorption remained stable. Thus, the ability of an adsorbent to retain the contaminant over time in soil may depend on adsorbent dose used. In our study, two doses were used, but very little or no difference was observed between doses. Further research will be required to determine whether PAH-ABC-soil complex is more stable with increased ABC dose and aging.

We investigated the role of DOC in PAH sorption to BC or ABC-amended soil. As hypothesized, higher levels of sorption were observed with sandy loam and sand, which had lower DOC concentrations in comparison with Haplic Arenosol. The result is in line with previous findings (Zhang et al. 2010; Kumari et al. 2014) where enhancement of PAH sorption in soil with BC was clearly observed in soils with low DOC and organic carbon contents. DOC associates strongly with BC surfaces in soil (Hale et al. 2011; Zhang et al. 2010), and SOM may block the pores of BC and reduce sorption of organic contaminants (Pignatello et al. 2006; Lian et al. 2015). After DOC was removed from soil there was enhanced PAH sorption to the Haplic Arenosol, not only to soil amended with BC, but also to control soil. This phenomenon in Haplic Arenosol could be explained by high native black carbon content compared with sandy loam and sand.

Also the third hypothesis of ours was supported by the evidence as both *A. praecox* and *P. velutina* increased sorption of pyrene to the Haplic Arenosol so that the bound fraction was approximately half of total pyrene. There seems to be a synergistic effect between BC and fungi in decreasing bioavailability and sorption of PAH, which was earlier confirmed

284 by García-Delgado et al. (2015) with low SOM soil, and our results further emphasize the role of SOM in the process.
285 White-rot and litter-decomposing fungi are known to produce oxidative enzymes, such as laccase and manganese
286 peroxidase, which are known to bind organic compounds to SOM through oxidative coupling (Berry and Boyd, 1984;
287 Bollag 1992; Held et al. 1997) as well as mineralize them (Steffen et al. 2002, Tuomela et al. 1999). However, we
288 measured no evolution of $^{14}\text{CO}_2$ during the entire incubation period. SOM influence has been observed to reduce
289 mineralization of pentachlorophenol (Tuomela et al. 1999) and the increase of aromatic rings and addition of BC probably
290 enhances the phenomenon as has been proved with ^{14}C -labelled synthetic lignin and phenanthrene (Tuomela et al. 2002,
291 Rhodes et al. 2008). If a contaminant is bound irreversibly to soil or BC, this can be considered as relevant remediation
292 action (Bollag 1992). In our earlier studies (Anasonye et al. 2014 and 2015) of soil system where a white-rot fungus was
293 grown (including *P. velutina*), manganese peroxidase was the main oxidative enzyme found with little or no laccase
294 activity, but García-Delgado et al. (2015) detected both laccase and manganese peroxidase from PAH-contaminated soil
295 with *Pleurotus ostreatus*. BC addition did not affect the enzyme production of the fungus. **In addition to BC and SOM,**
296 **bark acted as an adsorbent especially when organic matter content of soil was low, also previously reported (Olivella et**
297 **al. 2013, Winquist et al. 2014,). In addition to binding sites of the bark, bark extractives are capable of sorbing PAH**
298 **(Olivella et al. 2013).**

299

300 The results obtained from sterile soil with BC or ABC were compared with those obtained from non-sterile soil with
301 similar amendments in order to show the difference in roles of biotic versus abiotic processes in enhancement of pyrene
302 sorption in amended soil. The presence of native microbes in soil had higher impact on pyrene sorption to the amended-
303 Haplic Arenosol compared with the amended-sandy loam soil. Soil with high organic carbon tends to favor fungal
304 colonization over bacteria (Zhang et al. 2015), thus Haplic Arenosol, which is rich in lignocellulosic materials, was
305 probably more colonized with fungi capable of enhancing pyrene sorption as detected with *A. praecox* and *P. velutina*.

306 Stability of bound compound is very important as slow and continuous leaching to groundwater will pose a constant risk
307 to the environment (Bollag 1992). In our system, the soil was subjected to strong extraction procedure with organic solvent
308 to obtain the freely available fraction. The ^{14}C -pyrene fraction remaining in soil after strong chemical treatment is most
309 probably bound to adsorbents-SOM complex through covalent bonds, which are the most resistant to extraction and
310 degradation (Khan et al. 1978), and should remain in soil for a very long time.

311 **Conclusions**

312 We found that PAH can effectively be sorbed to carbonaceous soil amendments and thus be a promising tool for
313 remediating polluted sites. The effectiveness of biochar to sorb PAH can further be increased when activating the biochar
314 thus improving not only its BET surface area, but also its pore morphology and surface chemistry. The method was shown
315 to be even further enhanced if biochar addition is combined with treatment with basidiomycetous fungi, especially in high
316 SOM soils. Sorption of pyrene to BC-SOM complex enhanced by fungi could help reduce the leaching of pyrene from
317 soil to groundwater, but the results are yet to be validated under field conditions

318 **Acknowledgements**

319 The authors thank Jussi Heinonsalo, Kati Hakala and Kari Steffen for providing the experimental soils, Kaj-Roger Humm
320 for providing the guidance in working with labeled compounds. This research was funded by Maj and Tor Nessling
321 Foundation.

322

323 **References**

- 324 Abujabhah, I. S., Bound, S. A., Doyle, R., Bowman, J. P. (2016). Effects of biochar and compost amendments on soil
325 physico-chemical properties and the total community within a temperate agricultural soil. *Applied Soil Ecology*, 98, 243–
326 253.
- 327 Al Marzooqi, F., Yousef, L. F. (2017). Biological response of a sandy soil treated with biochar derived from a halophyte
328 (*Salicornia bigelovii*). *Applied Soil Ecology*, 114, 9–15.
- 329 Anasonye, F., Winquist, E., Räsänen, M., Kontro, J., Björklöf, K., Vasilyeva, G., Jørgensen, K. S., Steffen, K. T.,
330 Tuomela, M. (2015). Bioremediation of TNT contaminated soil with fungi under laboratory and pilot scale conditions.
331 *International Biodeterioration & Biodegradation*, 105, 7–12.
- 332 Anderson, C. R., Condron, L. M., Clough, T. J., Fiers, M., Stewart, A., Hill, R. A., Sherlock, R. R. (2011). Biochar
333 induced soil microbial community change: implications for biogeochemical cycling of carbon, nitrogen and phosphorus.
334 *Pedobiologia*, 54, 309–320.
- 335 Angst, T. E., Patterson, C. J., Reay, D. S., Anderson, P., Peshkur, T. A., Sohi, S. P. (2013). Biochar diminishes nitrous
336 oxide and nitrate leaching from diverse nutrient sources. *Journal of Environmental Quality*, 42, 672–682.
- 337 Anyika, C., Majid, Z. A., Ibrahim, Z., Zakaria, M. P., Yahya, A. (2014). The impact of biochars on sorption and
338 biodegradation of polycyclic aromatic hydrocarbons in soils—a review. *Environmental Science and Pollution Research*,
339 22, 3314–3341.
- 340 Beesley, L., Moreno-Jiménez, E., Gomez-Eyles, J. L., Harris, E., Robinson, B., Sizmur, T. (2011). A review of biochars’
341 potential role in the remediation, revegetation and restoration of contaminated soils. *Environmental pollution*, 159, 3269–
342 3282.
- 343 Berry D. F., Boyd, S. A. (1984). Oxidative coupling of phenols and anilines by peroxidase: structure-activity relationships.
344 *Soil Science Society of America Journal*, 48, 565–569.
- 345 Bollag, J. M. (1992). Decontaminating soil with enzymes. *Environmental Science & Technology*, 26, 1876–1881.
- 346 Case, S. D., McNamara, N. P., Reay, D. S., Whitaker, J. (2012). The effect of biochar addition on N₂O and CO₂ emissions
347 from a sandy loam soil—the role of soil aeration. *Soil Biology and Biochemistry*, 51, 125–134.

348 Chen, B., Yuan, M. (2011). Enhanced sorption of polycyclic aromatic hydrocarbons by soil amended with biochar.
349 *Journal of Soils and Sediments*, 11, 62–71.

350 Chen, S., Liao, C. (2006). Health risk assessment on human exposed to environmental polycyclic aromatic hydrocarbons
351 pollution sources. *Science of the Total Environment*, 366, 112–123.

352 Cheng, C., Lehmann, J. (2009). Ageing of black carbon along a temperature gradient. *Chemosphere*, 75, 1021–1027.

353 Cornelissen, G., Breedveld, G. D., Kalaitzidis, S., Christanis, K., Kibsgaard, A., Oen, A. M. (2006). Strong sorption of
354 native PAHs to pyrogenic and unburned carbonaceous geosorbents in sediments. *Environmental Science & Technology*,
355 40, 1197–1203.

356 Dai, Z., Hu, J., Zhang, L., Brookes, P. C., He, Y., Xu, J. (2016). Sensitive responders among bacterial and fungal
357 microbiome to pyrogenic organic matter (biochar) addition differed greatly between rhizosphere and bulk soils. *Scientific*
358 *Reports*, 6, 36101.

359 Dai Z, Hu J, Barberan A, Li Y, Brookes PC, He Y, Xu J (2017) Bacterial community composition associated with
360 pyrogenic organic matter (Biochar) varies with pyrolysis temperature and colonization environment. *Applied and*
361 *Environmental Science*, 2, 2 e00085-17.

362 de Andrés, J. M., Orjales, L., Narros, A., de la Fuente., María del Mar., Rodríguez M. E. (2013). Carbon dioxide
363 adsorption in chemically activated carbon from sewage sludge. *Journal of the Air & Waste Management Association*, 63,
364 557–564.

365 Dec, J., Haider, K., Bollag, J. (2001). Decarboxylation and demethoxylation of naturally occurring phenols during
366 coupling reactions and polymerization. *Soil Science*, 166, 660–671.

367 Deng, S., Zeng, D. (2017). Removal of phenanthrene in contaminated soil by combination of alfalfa, white-rot fungus
368 and earthworm. *Environmental Science and Pollution Research*, 24, 7565–7571.

369 Ding, Y., Liu, Y., Liu, S., Zhongwu, L., Tan, X., Huang, X., Zeng, G., Zhou, L., Zheng, B. (2016). Biochar to improve
370 soil fertility. *Agronomy for Sustainable Development*, 36, 1–18.

371 EBC, 2012. 'European biochar Certificate - Guidelines for a Sustainable Production of biochar. European biochar
372 Foundation (EBC), Arbaz, Switzerland. <http://www.europeanbiochar.org/en/download>. Version 6.2E of 04th February
373 2016. DOI: 10.13140/RG.2.1.4658.7043.

374 FAO-UNESCO. (1997). Soil map of the world. Revised legend, with corrections and updates. World Soil Resources
375 Report 60, Reprinted with updates as Technical paper 20, International Soil Reference and Information Centre,
376 Wageningen, 140 p.

377 Farrell, M., Kuhn, T. K., Macdonald, L. M., Maddern, T. M., Murphy, D. V., Hall, P. A., Singh, B. P., Baumann, k.,
378 Krull, E. S., Baldock, J. A. (2013). Microbial utilisation of biochar-derived carbon. *Science of the Total Environment*,
379 465, 288–297.

380 García-Delgado, C., Alfano-Barta, I., Eymar, E. (2015). Combination of biochar amendment and mycoremediation for
381 polycyclic aromatic hydrocarbons immobilization and biodegradation in creosote-contaminated soil. *Journal of*
382 *Hazardous Materials*, 285, 259–266.

383 Gibson, C., Berry, T. D., Wang, R., Spencer, J.A., Johnston, C. T., Jiang, Y., Bird, J. A., Filley, T. R. (2016). Weathering
384 of pyrogenic organic matter induces fungal oxidative enzyme response in single culture inoculation experiments. *Organic*
385 *Geochemistry*, 92, 32–41.

386 Grossman, J. M., O'Neill, B. E., Tsai, S. M., Liang, B., Neves, E., Lehmann, J., Thies, J.E. (2010). Amazonian anthrosols
387 support similar microbial communities that differ distinctly from those extant in adjacent, unmodified soils of the same
388 mineralogy. *Microbial Ecology*, 60, 192–205.

389 Hale, S., Hanley, K., Lehmann, J., Zimmerman, A., Cornelissen, G. (2011). Effects of chemical, biological, and physical
390 aging as well as soil addition on the sorption of pyrene to activated carbon and biochar. *Environmental Science &*
391 *Technology*, 45, 10445–10453.

392 Held, T., Draude, G., Schmidt, F., Brokamp, A., Reis, K. (1997). Enhanced humification as an in-situ bioremediation
393 technique for 2,4,6-trinitrotoluene (TNT) contaminated soils. *Environmental Science & Technology*, 18, 479–487.

394 Hilber, I., Blum, F., Leifeld, J., Schmidt, H. P., Bucheli, T. (2012) Quantitative determination of PAHs in biochar: a
395 prerequisite to ensure its quality and safe application. *Journal of Agricultural and Food Chemistry* 60, 3042–3050.

396 IBI (2015) Standardized product definition and product testing guidelines for biochar that is used in soil. International
397 biochar Initiative, p.15 [http://www. biocharinternational.org/sites/default/files/IBI_ Biochar_Standards_V2.1.pdf](http://www.biocharinternational.org/sites/default/files/IBI_Biochar_Standards_V2.1.pdf).
398 Accessed 18 December 2016.

399 Ilvesniemi, H., Giesler, R., van Hees, P., Magnusson, T., Melkerud, P. A. (2000). General description of the sampling
400 techniques and the sites investigated in the Fennoscandinavian podzolization project. *Geoderma* 94, 109–123.

401 Impellitteri, C. A., Lu, Y., Saxe, J. K., Allen, H. E., Peijnenburg, W. J. (2002). Correlation of the partitioning of dissolved
402 organic matter fractions with the desorption of Cd, Cu, Ni, Pb and Zn from 18 Dutch soils. *Environment International*,
403 28, 401–410.

404 Jin, H. (2010.) Characterization of microbial life colonizing biochar and biochar-amended soils. PhD Dissertation,
405 Cornell University, Ithaca, NY.

406 Jones, D., Rousk, J., Edwards-Jones, G., DeLuca, T., Murphy, D. (2012). Biochar-mediated changes in soil quality and
407 plant growth in a three year field trial. *Soil Biology and Biochemistry*, 45, 113–124.

408 Kästner, M., Nowak, K. M., Miltner, A., Stefan, Trapp, S., Schäffer, A. (2014). Classification and modelling of
409 nonextractable residue (NER) formation of xenobiotics in soil – a synthesis. *Critical Reviews in Environmental Science*
410 *and Technology*, 44, 2107–2171.

411 Khadem, A., Raiesi, F. (2017). Responses of microbial performance and community to corn biochar in calcareous sandy
412 and clayey soils. *Applied Soil Ecology*, 114, 16–27.

413 Khan, S. (1978). The interaction of organic matter with pesticides. In M. Schnitzer (Ed.), Soil organic matter: development
414 in soil science (pp. 137–171). Amsterdam; *Elsevier*.

415 Kolton, M., Meller, H. Y., Pasternak, Z., Graber, E. R., Elad, Y., Cytryn, E. (2011). Impact of biochar application to soil
416 on the root-associated bacterial community structure of fully developed greenhouse pepper plants. *Applied Environmental*
417 *Microbiology*, 77, 4924–4930.

418 Kumari, K., Moldrup, P., Paradelo, M., de Jonge, L.W. (2014). Phenanthrene sorption on biochar -amended soils:
419 Application rate, aging, and physicochemical properties of soil. *Water, Air, & Soil Pollution*, 225, 1–13.

420 Kurth, V., MacKenzie, M., DeLuca, T. (2006). Estimating charcoal content in forest mineral soils. *Geoderma*, 137, 135–
421 139.

422 Lalhruaitluanga, H., Prasad, M., Radha, K. (2011). Potential of chemically activated and raw charcoals of *Melocanna*
423 *baccifera* for removal of Ni (II) and Zn (II) from aqueous solutions. *Desalination*, 271, 301–308.

424 Lamichhane, S., Krishna, K., Sarukkalige. (2016). Polycyclic aromatic hydrocarbons (PAHs) removal by sorption: a
425 review. *Chemosphere*, 148, 336–353.

426 Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday, W. C., Crowley, D. (2011). Biochar effects on soil biota–
427 a review. *Soil Biology and Biochemistry*, 43, 1812–1836.

428 Lian, F., Sun, B., Chen, X., Zhu, L., Liu, Z., Xing, B. (2015). Effect of humic acid (HA) on sulfonamide sorption by
429 biochars. *Environmental pollution*, 204, 306–312.

430 Liedekerke, M., Prokop, G., Rabl-Berger, S., Kibblewhite and G.Louwagie. (2014). Progress in the management of
431 contaminated sites in Europe. JRC Reference Reports, Joint Research Centre, Report EUR 26376 EN, European
432 Commission. <http://www.eea.europa.eu/data-and-maps/indicators/progress-in-management-of-contaminated-sites-3/joint-research-centre-2014-progress>. Accessed 06 September 2016.

433

434 Liu, P., Liu, W., Jiang, H., Chen, J., Li, W., Yu, H. (2012). Modification of bio-char derived from fast pyrolysis of biomass
435 and its application in removal of tetracycline from aqueous solution. *Bioresource Technology*, 121, 235–240.

436 Macleod, C. J., Semple, K.T. (2002). The adaptation of two similar soils to pyrene catabolism. *Environmental pollution*,
437 119, 357–364.

438 Major, J., Rondon, M., Molina, D., Riha, S. J., Lehmann, J. (2010). Maize yield and nutrition during 4 years after biochar
439 application to a Colombian savanna oxisol. *Plant Science*, 333, 117–128.

440 Martin, S. M., Kookana, R. S., Van Zwieten, L., Krull, E. (2012). Marked changes in herbicide sorption–desorption upon
441 ageing of biochars in soil. *Journal of Hazardous Materials*, 231, 70–78.

442 Mitchell, P. J., Dalley, T. S., Helleur, R. J. (2013). Preliminary laboratory production and characterization of biochars
443 from lignocellulosic municipal waste. *Journal of Analytical and Applied Pyrolysis*, 99, 71–78.

444 Mitchell, P. J., Simpson, A. J., Soong, R., Schurman, J. C., Thomas, S. C., Simpson, M. J. (2016) Biochar amendment
 445 and phosphorus altered forest soil microbial community and native soil organic matter molecular composition.
 446 *Biogeochemistry*, 130, 227–245. Murphy, B.L., Brown, J. (2005). Environmental forensics aspects of PAHs from wood
 447 treatment with creosote compounds. *Environmental Forensics*, 6, 151–159.

448 Nielsen, S., Minchin, T., Kimber, S., van Zwieten, L., Gilbert, J., Munroe, P., Joseph, S., Thomas, T. (2014). Comparative
 449 analysis of the microbial communities in agricultural soil amended with enhanced biochars or traditional fertilisers.
 450 *Agriculture, Ecosystems & Environment*, 191, 73–82.

451 Noyce, G. L., Winsborough, C., Fulthorpe, R., Basiliko, N. (2016). The microbiomes and metagenomes of forest biochars.
 452 *Scientific Reports*, 6, 26425.

453 O'Neill, B., Grossman, J., Tsai, M., Gomes, J., Lehmann, J., Peterson, J., Neves, E., Thies, J. E. (2009). Bacterial
 454 community composition in Brazilian anthrosols and adjacent soils characterized using culturing and molecular
 455 identification. *Microbial Ecology*, 58, 23–35.

456 Ogbonnaya, U., Oyelami, A., Matthews, J., Adebisi, O., Semple, K. T. (2014). Influence of wood biochar on phenanthrene
 457 catabolism in soils. *Environments*, 1, 60–74.

458 Olivella, Costa, À., Fernández, I., Cano, L., Jové, P., Oliveras, A. (2013). Role of chemical components of cork on
 459 sorption of aqueous polycyclic aromatic hydrocarbons. *International Journal of Environmental Research*, 1, 225–234.

460 Pan, F., Li, Y., Chapman, S. T., Khan, S., Yao, H. (2016). Microbial utilization of rice straw and its derived biochar in a
 461 paddy Soil. *Science of the Total Environment*, 559, 15–23.

462 Park, J., Hung, I., Gan, Z., Rojas, O. J., Lim, K. H., Park, S. (2013). Activated carbon from biochar: Influence of its
 463 physicochemical properties on the sorption characteristics of phenanthrene. *Bioresource Technology*, 149, 383–389.

464 Pietikäinen, J., Kiikkilä, O., Fritze, H. (2000). Charcoal as a habitat for microbes and its effect on the microbial community
 465 of the underlying humus. *Oikos*, 89, 231–242.

466 Pignatello, J. J., Kwon, S., Lu, Y. (2006). Effect of natural organic substances on the surface and adsorptive properties of
 467 environmental black carbon (char): attenuation of surface activity by humic and fulvic acids. *Environmental Science &*
 468 *Technology*, 40, 7757–7763.

469 Prayogo, C., Jones, J. E., Bending, G. D. (2013). Impact of biochar on mineralisation of C and N from soil and willow
 470 litter and its relationship with microbial community biomass and structure. *Biology and Fertility of Soils*, 50, 695–702.

471 Quilliam, R.S., Glanville, H. C., Wade, S. C., Jones, D. L. (2013a). Life in the charosphere – does biochar in agricultural
 472 soil provide a significant habitat for microorganisms? *Soil Biology and Biochemistry*, 69, 287–293.

473 Quilliam, R. S., Rangecroft, S., Emmett, B. A., Deluca, T.H., Jones, D. L. (2013b). Is biochar a source or sink for
 474 polycyclic aromatic hydrocarbon (PAH) compounds in agricultural soils? *GCB Bioenergy*, 5, 96–103.

475 Rhodes, A., Carlin, A., Semple, K. T. (2008). Impact of Black Carbon in the Extraction and Mineralization of
 476 Phenanthrene in Soil. *Science of the Total Environment*, 42, 740–745.

477 Rumpel, C., Alexis, M., Chabbi, A., Chaplot, V., Rasse, D. P., Valentin, C., Mariotti, A. (2006). Black carbon contribution
478 to soil organic matter composition in tropical sloping land under slash and burn agriculture. *Geoderma* 130, 35–46.

479 Smith, P. (2016). Soil carbon sequestration and biochar as negative emission technologies. *Global Change Biology*, 22,
480 1315–1324.

481 Soil Survey Staff. (1998). Keys to Soil Taxonomy. United States Department of Agriculture/Natural Resources
482 Conservation Service.

483 Steffen, K., Hatakka, A., Hofrichter, M. (2002). Removal and mineralization of polycyclic aromatic hydrocarbons by
484 litter-decomposing basidiomycetous fungi. *Applied Microbiology and Biotechnology*, 60, 212–217.

485 Tammeorg, P., Parviainen, T., Nuutinen, V., Simojoki, A., Vaara, E., Helenius, J. (2014a). Effects of biochar on
486 earthworms in arable soil: avoidance test and field trial in boreal loamy sand. *Agriculture, Ecosystems & Environment*,
487 191, 150–157.

488 Tammeorg, P., Simojoki, A., Mäkelä, P., Stoddard, F. L., Alakukku, L., Helenius, J. (2014b). Short-term effects of biochar
489 on soil properties and wheat yield formation with meat bone meal and inorganic fertiliser on a boreal loamy sand.
490 *Agriculture, Ecosystems & Environment*, 191, 108–116.

491 Tuomela, M., Lyytikäinen, M., Oivanen, P., Hatakka, A. (1999). Mineralization and conversion of pentachlorophenol
492 (PCP) in soil inoculated with the white-rot fungi *Trametes versicolor*. *Soil Biology and Biochemistry*, 31, 65–74.

493 Tuomela, M., Oivanen, P., Hatakka, A. (2002). Degradation of synthetic ¹⁴C-lignin by various white-rot fungi in soil. *Soil*
494 *Soil Biology and Biochemistry*, 34, 1613–1620.

495 Valentín, L., Oesch-Kuisma, H., Steffen, K. T., Kähkönen, M. A., Hatakka, A., Tuomela, M. (2013). Mycoremediation
496 of wood and soil from an old sawmill area contaminated for decades. *Journal of Hazardous Materials*, 260, 668–675.

497 Watzinger, A., Feichtmair, S., Kitzler, B., Zehetner, F., Kloss, S., Wimmer, B., Zechmeister-Boltenstern, S., Soja, G.
498 (2014). Soil microbial communities responded to biochar application in temperate soils and slowly metabolized ¹³C-
499 labelled biochar as revealed by ¹³C PLFA analyses: results from a short-term incubation and pot experiment. *European*
500 *Journal of Soil Science*, 65, 40–51.

501 Warnock, D. D., Lehmann, J., Kuypers, T. W., Rillig, M. C. (2007). Mycorrhizal responses to biochar in soil—concepts
502 and mechanisms. *Plant Soil*, 300, 9–20.

503 Winquist, E., Björklöf, K., Schultz, E., Räsänen, M., Salonen, K., Anasonye, F., Cajthaml, T., Steffen, K. T., Jørgensen,
504 K. S., Tuomela, M. (2014). Bioremediation of PAH-contaminated soil with fungi—From laboratory to field scale.
505 *International Biodeterioration & Biodegradation*, 86, 238–247.

506 Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., Joseph, S. (2010). Sustainable biochar to mitigate global
507 climate change. *Nature Communications*, 1, 56.

508 Zhang, Q., Zhou, W., Liang, G. Q., Sun, J. W., Wang, X. B., He, P. (2015). Distribution of soil nutrients, extracellular
509 enzyme activities and microbial communities across particle-size fractions in a long-term fertilizer experiment. *Applied*
510 *Soil Ecology*, 94, 59–71.

511 Zhang, H., Lin, K., Wang, H., Gan, J. (2010). Effect of *Pinus radiata* derived biochar on soil sorption and desorption of
512 phenanthrene. *Environmental pollution*, 158, 2821–2825.

513 Zhu, X., Chen, B., Zhu, L., Xing, B. (2017). Effects and mechanisms of biochar-microbe interactions in soil improvement
514 and pollution remediation: a review. *Environmental pollution*, 227, 98–115.

515

516 Figure caption

517 **Figure 1** Sorption parameter (K_d) ($L\ kg^{-1}$) of ^{14}C -pyrene in soil with (on left) and without DOC (on right) with or without
518 1% (w/w) BC or ABC. Dissolved organic matter (DOC) was extracted from the soil with deionized water three times

519